Proof of a supercongruence conjectured by Z.-H. Sun

Victor J. W. Guo

Department of Mathematics, Shanghai Key Laboratory of PMMP, East China Normal University, 500 Dongchuan Rd., Shanghai 200241, People's Republic of China

jwguo@math.ecnu.edu.cn, http://math.ecnu.edu.cn/~jwguo

arXiv:1404.6978v1 [math.NT] 28 Apr 2014

Abstract. The Franel numbers are defined by $f_n=\sum_{k=0}^n\binom{n}{k}^3$. Motivated by the recent work of Z.-W. Sun on Franel numbers, we prove that

$$\sum_{k=0}^{n-1}(3k+1)(-16)^{n-k-1}\binom{2k}{k}f_k\equiv 0 \pmod{n\binom{2n}{n}},$$

$$\sum_{k=0}^{p-1}\frac{3k+1}{(-16)^k}\binom{2k}{k}f_k\equiv p(-1)^{\frac{p-1}{2}} \pmod{p^3}.$$

where $n>1$ and p is an odd prime. The second congruence modulo p^2 confirms a recent conjecture of Z.-H. Sun. We also show that, if p is a prime of the form $4k+3$, then

$$\sum_{k=0}^{p-1}\frac{\binom{2k}{k}f_k}{(-16)^k}\equiv 0 \pmod{p},$$

which confirms a special case of another conjecture of Z.-H. Sun.

Keywords: Franel numbers, binomial coefficients, multinomial coefficients, congruences

AMS Subject Classifications: 11A07, 11B65, 05A10, 05A19

1 Introduction

The numbers f_n, defined by

$$f_n=\sum_{k=0}^n\binom{n}{k}^3,$$

were first studied by Franel [3, 4], who obtained the following recurrence relation:

$$(n+1)^2f_{n+1}=(7n^2+7n+2)f_n+8n^2f_{n-1},\ n=1,2,\ldots.$$

MacMahon [8, p. 122] gave the following identity related to Franel numbers:

$$\sum_{k=0}^n\binom{n}{k}^3x^k=\sum_{k=0}^n\binom{n+k}{3k}\binom{3k}{2k}\binom{2k}{k}x^k(1+x)^{n-2k} \tag{1.1}$$

(see also Foata [2] or Riordan [11, p. 41]). Strehl [12] gave another expression for Franel numbers:

$$f_n = \sum_{k=0}^{n} \binom{n}{k}^2 \binom{2k}{n}.$$

Jarvis and Verrill [6] obtained the following congruence:

$$f_n \equiv (-8)^n f_{p-1-n} \pmod{p},$$

where p is a prime and $0 \le n \le p-1$. Recently, Z.-W. Sun [17, 18] proved many interesting congruences involving Franel numbers. On the other hand, during his working on congruences for Legendre polynomials [13–15], Z.-H. Sun [16] proposed a lot of conjectures on supercongruences concerning Franel numbers , such as

Conjecture 1.1. (see [16, Conjecture 4.23])*Let $p > 3$ be a prime. Then*

$$\sum_{k=0}^{p-1} \frac{3k+1}{(-16)^k} \binom{2k}{k} f_k \equiv p(-1)^{\frac{p-1}{2}} \pmod{p^2}.$$

Conjecture 1.2. [16, Conjecture 4.14]*Let p be an odd prime. Then*

$$\sum_{k=0}^{p-1} \frac{\binom{2k}{k} f_k}{(-16)^k} \equiv \begin{cases} 4x^2 - 2p \pmod{p^2}, & \text{if } p = x^2 + y^2 \equiv 1 \bmod 12 \text{ with } 6 \mid y, \\ 2p - 4x^2 \pmod{p^2}, & \text{if } p = x^2 + y^2 \equiv 1 \bmod 12 \text{ with } 6 \mid x-3, \\ 4\left(\frac{xy}{3}\right) xy \pmod{p^2}, & \text{if } p = x^2 + y^2 \equiv 5 \bmod 12, \\ 0 \pmod{p^2}, & \text{if } p \equiv 3 \bmod 4, \end{cases}$$

where $\left(\frac{a}{3}\right)$ is the Legendre symbol.

In this paper, we shall prove the following results.

Theorem 1.3. *Let $n > 1$ be a positive integer. Then*

$$\sum_{k=0}^{n-1} (3k+1)(-16)^{n-k-1} \binom{2k}{k} f_k \equiv 0 \pmod{n\binom{2n}{n}}. \tag{1.2}$$

Theorem 1.4. *Let p be an odd prime. Then*

$$\sum_{k=0}^{p-1} \frac{3k+1}{(-16)^k} \binom{2k}{k} f_k \equiv p(-1)^{\frac{p-1}{2}} \pmod{p^3}. \tag{1.3}$$

Theorem 1.5. *Let p be a prime of the form $4k+3$. Then*

$$\sum_{k=0}^{p-1} \frac{\binom{2k}{k} f_k}{(-16)^k} \equiv 0 \pmod{p}. \tag{1.4}$$

It is obvious that the congruence (1.3) modulo p^2 confirms Conjecture 1.1, while (1.4) is a special case of Conjecture 1.2.

2 Proof of Theorem 1.3

We need the following identity due to Z.-W. Sun [17, (2.3)]:

$$f_n=\sum_{k=0}^{n}\binom{n+2k}{3k}\binom{3k}{k}\binom{2k}{k}(-4)^{n-k}, \tag{2.1}$$

which can be proved by the Zeilberger algorithm (see [7, 10]). Moreover, by induction on n, we can easily prove that, for all $0\leqslant k\leqslant n$,

$$\sum_{m=k}^{n-1}(3m+1)(-16)^{n-m-1}\binom{2m}{m}\binom{m+2k}{3k}(-4)^{m-k}=\binom{2n}{n}\binom{n+2k}{3k}\frac{n(k-n)(-4)^{n-k}}{8(2k+1)}. \tag{2.2}$$

In fact, when $n=k$, each side of (2.2) equals 0. Suppose that (2.2) is true for n. Then

$$\begin{aligned}
&\sum_{m=k}^{n}(3m+1)(-16)^{n-m}\binom{2m}{m}\binom{m+2k}{3k}(-4)^{m-k}\\
&=(3n+1)\binom{2n}{n}\binom{n+2k}{3k}(-4)^{n-k}\\
&\quad+\sum_{m=k}^{n-1}(3m+1)(-16)^{n-m}\binom{2m}{m}\binom{m+2k}{3k}(-4)^{m-k}\\
&=(3n+1)\binom{2n}{n}\binom{n+2k}{3k}(-4)^{n-k}-2\binom{2n}{n}\binom{n+2k}{3k}\frac{n(k-n)(-4)^{n-k}}{(2k+1)}\\
&=\binom{2n+2}{n+1}\binom{n+2k+1}{3k}\frac{(n+1)(k-n-1)(-4)^{n-k+1}}{8(2k+1)}.
\end{aligned}$$

Namely, the identity (2.2) holds for $n+1$.

By (2.1) and (2.2), we have

$$\begin{aligned}
&\sum_{m=0}^{n-1}(3m+1)(-16)^{n-m-1}\binom{2m}{m}f_m\\
&\quad=\sum_{m=0}^{n-1}(3m+1)(-16)^{n-m-1}\binom{2m}{m}\sum_{k=0}^{m}\binom{m+2k}{3k}\binom{3k}{k}\binom{2k}{k}(-4)^{m-k}.\\
&\quad=\sum_{k=0}^{n-1}\binom{2n}{n}\binom{n+2k}{3k}\binom{3k}{k}\binom{2k}{k}\frac{n(k-n)(-4)^{n-k}}{8(2k+1)}.
\end{aligned} \tag{2.3}$$

Note that, for $n\geqslant 2$ and $0\leqslant k<n$, both

$$\frac{1}{2k+1}\binom{3k}{k}=\binom{3k}{k}-2\binom{3k}{k-1}$$

and $\binom{2k}{k}\frac{(-4)^{n-k}}{8}$ are integers. From (2.3) we deduce that

$$\frac{1}{n\binom{2n}{n}}\sum_{k=0}^{n-1}(3k+1)(-16)^{n-k-1}\binom{2k}{k}f_k=\sum_{k=0}^{n-1}\binom{n+2k}{3k}\binom{3k}{k}\binom{2k}{k}\frac{(k-n)(-4)^{n-k}}{8(2k+1)} \quad (2.4)$$

is an integer. This proves (1.2).

3 Proof of Theorem 1.4

It follows from (2.4) that

$$\frac{1}{p}\sum_{k=0}^{p-1}\frac{3k+1}{(-16)^k}\binom{2k}{k}f_k=-4^{1-p}\binom{2p-1}{p-1}\sum_{k=0}^{p-1}\binom{p+2k}{3k}\binom{3k}{k}\binom{2k}{k}\frac{k-p}{(2k+1)(-4)^k}. \quad (3.1)$$

Note that Babbage [1] proved the following congruence:

$$\binom{2p-1}{p-1}\equiv 1 \pmod{p^2}. \quad (3.2)$$

Moreover, we have

$$\binom{p+2k}{3k}\binom{3k}{k}=\frac{(p+2k)!}{(2k)!k!(p-k)!}\equiv\begin{cases}(-1)^{k-1}\frac{p}{k}, & \text{if } 1\leqslant k<\frac{p-1}{2},\\ (-1)^{k-1}\frac{2p}{k}, & \text{if } \frac{p+1}{2}\leqslant k<p\end{cases}\pmod{p^2}, \quad (3.3)$$

and, when $k=\frac{p-1}{2}$,

$$\binom{p+2k}{3k}\binom{3k}{k}\binom{2k}{k}\frac{k-p}{2k+1}=-\binom{2p-1}{p-1}\binom{p-1}{\frac{p-1}{2}}^2\equiv -16^{p-1} \pmod{p^2}, \quad (3.4)$$

where we have used Babbage's congruence (3.2) and Morley's congruence [9]:

$$\binom{p-1}{\frac{p-1}{2}}\equiv(-1)^{\frac{p-1}{2}}4^{p-1} \pmod{p^3}.$$

Substituting (3.2)–(3.4) into (3.1), and observing that $\binom{2k}{k}\equiv 0 \pmod{p}$ for $\frac{p-1}{2}<k<p$, we obtain

$$\begin{aligned}\frac{1}{p}\sum_{k=0}^{p-1}\frac{3k+1}{(-16)^k}\binom{2k}{k}f_k&\equiv p4^{1-p}+(-4)^{\frac{p-1}{2}}+4^{1-p}\sum_{k=1}^{\frac{p-3}{2}}\binom{2k}{k}\frac{p-p^2/k}{(2k+1)4^k}\\ &\equiv(-4)^{\frac{p-1}{2}}+4^{1-p}\sum_{k=0}^{\frac{p-3}{2}}\binom{2k}{k}\frac{p}{(2k+1)4^k} \pmod{p^2}. \quad (3.5)\end{aligned}$$

Since

$$\binom{2k}{k}4^{-k}\equiv(-1)^k\binom{\frac{p-1}{2}}{k}\pmod p, \tag{3.6}$$

we may rewrite (3.5) as

$$\frac{1}{p}\sum_{k=0}^{p-1}\frac{3k+1}{(-16)^k}\binom{2k}{k}f_k\equiv(-4)^{\frac{p-1}{2}}+4^{1-p}\sum_{k=0}^{\frac{p-3}{2}}(-1)^k\binom{\frac{p-1}{2}}{k}\frac{p}{2k+1}\pmod{p^2}, \tag{3.7}$$

Applying the famous identity

$$\sum_{k=0}^{n}(-1)^k\binom{n}{k}\frac{1}{x+k}=\frac{n!}{x(x+1)\cdots(x+n)}$$

with $x=\frac{1}{2}$ and $n=\frac{p-1}{2}$, we may simplify (3.7) as

$$\begin{aligned}\frac{1}{p}\sum_{k=0}^{p-1}\frac{3k+1}{(-16)^k}\binom{2k}{k}f_k&\equiv(-4)^{\frac{p-1}{2}}+2^{1-p}\binom{p-1}{\frac{p-1}{2}}^{-1}-(-1)^{\frac{p-1}{2}}4^{1-p}\\&\equiv(-1)^{\frac{p-1}{2}}\left(2^{p-1}+8^{1-p}-4^{1-p}\right)\pmod{p^2}.\end{aligned}$$

By Fermat's little theorem, we have $2^{p-1}-1\equiv 0\pmod p$ and so

$$2^{p-1}+8^{1-p}-4^{1-p}-1=(2^{p-1}-1)^2(2^{1-p}+4^{1-p}+8^{1-p})\equiv 0\pmod{p^2}.$$

This completes the proof. □

4 Proof of Theorem 1.5

We first give a binomial coefficient identity.

Lemma 4.1. *Let n and k be nonnegative integers with $k\leqslant n$. Then*

$$\sum_{m=k}^{n}\binom{n}{m}\binom{m+2k}{3k}(-1)^{m-k}=\binom{2k}{n-k}(-1)^{n-k}. \tag{4.1}$$

Proof. Consider the generating function of the left-hand side of (4.1). By the binomial theorem, we have

$$\begin{aligned}\sum_{n=k}^{\infty}x^n\sum_{m=k}^{n}\binom{n}{m}\binom{m+2k}{3k}(-1)^{m-k}&=\sum_{m=k}^{\infty}\binom{m+2k}{3k}(-1)^{m-k}\sum_{n=m}^{\infty}\binom{n}{m}x^n\\&=\sum_{m=k}^{\infty}\binom{m+2k}{3k}\frac{(-1)^{m-k}x^m}{(1-x)^{m+1}}\\&=\frac{x^k}{(1-x)^{x+1}}\left(1+\frac{x}{1-x}\right)^{-3k-1}\\&=x^k(1-x)^{2k},\end{aligned}$$

which is clearly the generating function of the right-hand side of (4.1). □

By (3.6) and (2.1), we have

$$\sum_{k=0}^{p-1}\frac{\binom{2k}{k}f_k}{(-16)^k}\equiv\sum_{m=0}^{p-1}\frac{\binom{\frac{p-1}{2}}{m}f_m}{4^m}$$
$$=\sum_{m=0}^{p-1}\frac{\binom{\frac{p-1}{2}}{m}}{4^m}\sum_{k=0}^{m}\binom{m+2k}{3k}\binom{3k}{k}\binom{2k}{k}(-4)^{m-k}\pmod p. \tag{4.2}$$

Exchanging the summation order in the right-hand side of (4.2), and applying (4.1) with $n=\frac{p-1}{2}$, we obtain

$$\sum_{k=0}^{p-1}\frac{\binom{2k}{k}f_k}{(-16)^k}\equiv\sum_{k=0}^{\frac{p-1}{2}}\frac{(-1)^{\frac{p-1}{2}-k}}{4^k}\binom{3k}{k}\binom{2k}{k}\binom{2k}{\frac{p-1}{2}-k}\pmod p. \tag{4.3}$$

It is easy to see that

$$\binom{2k}{\frac{p-1}{2}-k}\equiv\binom{\frac{3(p-1)}{2}-3k}{\frac{p-1}{2}-k}(-1)^{\frac{p-1}{2}-k}\pmod p. \tag{4.4}$$

Thus, by (3.6) and (4.4), we may rewrite (4.3) as

$$\sum_{k=0}^{p-1}\frac{\binom{2k}{k}f_k}{(-16)^k}\equiv\sum_{k=0}^{\frac{p-1}{2}}(-1)^k\binom{\frac{p-1}{2}}{k}\binom{3k}{k}\binom{\frac{3(p-1)}{2}-3k}{\frac{p-1}{2}-k}\pmod p. \tag{4.5}$$

If p is a prime of the form $4k+3$, then $\frac{p-1}{2}$ is odd, and by the symmetry of binomial coefficients, the right-hand side of (4.5) equals 0. This completes the proof.

5 Concluding remarks and open problems

Inspired by Conjectures 4.22 and 4.23 in [16] and Theorem 1.3 in this paper, we propose the following two conjectures.

Conjecture 5.1. *Let $n>1$ be a positive integer and let $(a,b,c)\in\{(9,4,5),(5,2,16),(9,2,50),(5,1,96),(6,1,320),(90,13,896),(102,11,10400)\}$. Then*

$$\sum_{k=0}^{n-1}(ak+b)c^{n-k-1}\binom{2k}{k}f_k\equiv 0\pmod{n\binom{2n}{n}}.$$

Conjecture 5.2. *Let $n>1$ be a positive integer and let $(a,b,c)\in\{(15,4,-49),(9,2,-112),(99,17,-400),(855,109,-2704),(585,58,-24304)\}$. Then*

$$\sum_{k=0}^{n-1}(ak+b)c^{n-k-1}\binom{2k}{k}f_k\equiv 0\pmod{n\binom{2n}{n}}.$$

We have verified Conjectures 5.1 and 5.2 for n up to 500 via Maple.

We also make the third conjecture on Franel numbers as follows.

Conjecture 5.3. *Let m, n be positive integers and let $a_1, \ldots, a_m$ be integers. Then*

$$\sum_{k=0}^{n-1}(3k+2)(-1)^{(m-1)k}f_k\prod_{i=1}^{m}\binom{a_in-1}{k}\binom{a_in+k}{k}\equiv 0 \pmod{n^2}, \tag{5.1}$$

$$\sum_{k=0}^{n-1}(9k^2+5k)(-1)^{(m-1)k}f_k\prod_{i=1}^{m}\binom{a_in-1}{k}\binom{a_in+k}{k}\equiv 0 \pmod{n^2}. \tag{5.2}$$

Note that, for any prime p and nonnegative integer $k \leqslant p-1$, there holds

$$\binom{a_ip-1}{k}\binom{a_ip+k}{k}=\prod_{j=1}^{k}\frac{(a_ip-j)(a_ip+j)}{j^2}\equiv(-1)^k \pmod{p^2}.$$

Therefore, by [17, Theorem 1.1], the congruences (5.1) and (5.2) are true for any prime n. On the other hand, the author [5] has proved that

$$\sum_{k=0}^{n-1}(3k+2)(-1)^k f_k\equiv 0 \pmod{2n^2},$$

which was conjectured by Z.-W. Sun (see [18, Conjecture 1.3]); and for $m=0$ and $n>1$, the congruence (5.2) can be further strengthened as

$$\sum_{k=0}^{n-1}(9k^2+5k)(-1)^k f_k\equiv 0 \pmod{n^2(n-1)},$$

which was also conjectured by Z.-W. Sun (see Conjecture 5.3 in a previous version of [18]: `http://arxiv.org/pdf/1112.1034v10.pdf`).

Acknowledgments. This work was partially supported by the Fundamental Research Funds for the Central Universities and the National Natural Science Foundation of China (grant 11371144).

References

[1] C. Babbage, Demonstration of a theorem relating to prime numbers, Edinburgh Philos. J. 1 (1819), 46–49.

[2] D. Foata, Etude algébrique de certains problèmes d'analyse combinatoire et du calcul des probabilités, Publ. Inst. Statist. Univ. Paris 14 (1965), 81–241.

[3] J. Franel, On a question of Laisant, L'intermédiaire des mathématiciens 1 (1894), 45–47.

[4] J. Franel, On a question of J. Franel, L'intermédiaire des mathématiciens 2 (1895), 33–35.

[5] V.J.W. Guo, Proof of two conjectures of Sun on congruences for Franel numbers, Integral Transforms Spec. Funct. 24 (2013), 532–539.

[6] F. Jarvis and H. A. Verrill, Supercongruences for the Catalan-Larcombe-French numbers, Ramanujan J. 22 (2010), 171–186.

[7] W. Koepf, Hypergeometric Summation, an Algorithmic Approach to Summation and Special Function Identities, Friedr. Vieweg & Sohn, Braunschweig, 1998.

[8] P. A. MacMachon, Combinatorial Analysis, Vol. 1, Cambridge University Press, London, 1915.

[9] F. Morley, Note on the congruence $2^{4n} \equiv (-1)^n (2n)!/(n!)^2$, where $2n+1$ is a prime, Ann. Math. 9 (1894/95), 168–170.

[10] M. Petkovšek, H. S. Wilf, and D. Zeilberger, $A = B$, A K Peters, Ltd., Wellesley, MA, 1996.

[11] J. Riordan, Combinatorial Identities, J. Wiley, New York, 1979.

[12] V. Strehl, Binomial sums and identities, Maple Technical Newsletter 10 (1993), 37–49.

[13] Z.-H. Sun, Congruences concerning Legendre polynomials, Proc. Amer. Math. Soc. 139 (2011), 1915–1929.

[14] Z.-H. Sun, Congruences concerning Legendre polynomials II, J. Number Theory 133 (2013), 1950–1976.

[15] Z.-H. Sun, Congruences concerning Legendre polynomials III, Int. J. Number Theory 9 (2013), 965–999.

[16] Z.-H. Sun, Some conjectures on congruences, preprint, 2013, arXiv:1103.5384v5.

[17] Z.-W. Sun, Connections between $p = x^2 + 3y^2$ and Franel numbers, J. Number Theory 133 (2013), 2914–2928.

[18] Z.-W. Sun, Congruences for Franel numbers, Adv. Appl. Math. 51 (2013), 524–535.